

**DISTREPTAZA**  
Streptokinase + Streptodornase  
15000 IU + 1250 IU  
Rectal suppositories

**Read the package leaflet before use. It contains important patient information.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it to others.
- It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What is Distreptaza and what it is used for
2. Before you receive Distreptaza
3. How Distreptaza is given
4. Possible side effects
5. How to store Distreptaza
6. Further information

1. What is Distreptaza and what it is used for  
Distreptaza comes in the form of rectal suppositories containing two active substances: streptokinase and streptodornase.

It dissolves blood clots and pus, which facilitates better access of antibiotics, chemotherapeutics and antibodies to the focus of inflammation.

It is used, to an excellent effect, in the following diseases:

- chronic adnexitis,
- infiltrative/postoperative
- lesions, endometritis,
- acute and chronic inflammatory conditions of haemorrhoidal
- varices, perirectal abscesses and fistulas with extensive
-

inflammatory infiltration,  
suppurating cysts of coccygeal bone region (coccygeal bone, also referred to as the tail bone, is located in the lower part of the spine, below the sacral bone).

## **2. Before you receive**

### **Distreptaza**

#### **Do not receive Distreptaza if:**

Contraindications for use may include: hypersensitivity to any of product ingredients, bleeding diatheses, wounds covered with fresh crust or surgical sutures within the administration area.

#### Warnings and precautions

The medicine should not come into contact with wounds covered with newly formed crust or surgical sutures, as this may lead to suture loosening and secondary bleeding.

#### Taking other medicines

Distreptaza should not be received together with antithrombotic drugs, as this may lead to local bleeding.

### **Pregnancy and breastfeeding**

Distreptaza should not be given to women who are pregnant or breastfeeding.

### **Driving and using machines**

Distreptaza produces no effect on the ability to drive and use machines.

## **3. How Distreptaza is given**

Always take Distreptaza according to doctor's prescription.

After removing from a blister pack cavity, the suppository should be inserted deep into the rectum.

Dosage depends on the type and acuteness of the inflammatory state.

In serious conditions, the following quantities are administered respectively:

3 times a day 1 suppository for the initial 3 days  
2 times a day 1 suppository for 3 consecutive days  
once daily 1 suppository for 3 consecutive days  
In less serious conditions:

2 times a day 1 suppository for 3 days  
once daily 1 suppository for 4 consecutive days

or

2 times a day 1 suppository for 2 days

Average number of suppositories during a cycle of treatment is 8-18.

Average treatment cycle is 7-10 days.

#### 4. Possible side effects

Like all medicines, Distreptaza can cause side effects, although not everybody gets them.

In general, it is well-tolerated.

Rarely, allergic reactions, increased body temperature and bleeding diathesis may occur.

If you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### 5. How to store Distreptaza

Keep out of the reach and sight of children.

Store in temperature 2°C - 8°C (refrigerated). Do not freeze.

Do not use after expiry date stated on packaging.

Unused medicinal product or any waste materials derived from it must be disposed of properly.

Lot number

Expiry date

## **6. Packaging content and further information**

What Distreptaza contains

One suppository contains:

Streptokinase 15000 IU

Streptodornase 1250 IU

Excipients: Paraffin oil, Witepsol H15

### **What Distreptaza looks like and contents of the pack**

#### **What Distreptaza looks like**

Distreptaza has the form of a white to cream coloured torpedo-shaped suppository.

#### **Packaging**

1 blister pack containing 6 suppositories 2 blister packs, each containing 5 suppositories Blister pack of PVC/PE in a cardboard box.

#### **Marketing Authorisation Holder and Manufacturer:**

„BIOMED-LUBLIN” Wytwórnia Surowic i Szczepionek Spółka Akcyjna

20-029 Lublin,

ul. Uniwersytecka 10

Phone: + 48 (81) 533 82 21

e-mail: [biomed@biomed.lublin.pl](mailto:biomed@biomed.lublin.pl)

This leaflet was approved in: **2011-11-28**